

## THE GENERAL RELATIVITY WITH CONFORMAL UNITS

*V. N. Pervushin*<sup>1</sup>, *A. B. Arbuzov*<sup>1,2</sup>, *B. M. Barbashov*<sup>1</sup>,  
*R. G. Nazmitdinov*<sup>1,3</sup>, *A. Borowiec*<sup>4</sup>, *K. N. Pichugin*<sup>5</sup>,  
*A. F. Zakharov*<sup>6</sup>

<sup>1</sup> Joint Institute for Nuclear Research, Dubna

<sup>2</sup> Department of Higher Mathematics, University of Dubna, Dubna, Russia

<sup>3</sup> Department de Física, Universitat de les Illes Balears, Palma de Mallorca, Spain

<sup>4</sup> Institute of Theoretical Physics, University of Wrocław, Wrocław, Poland

<sup>5</sup> Kirensky Institute of Physics, Krasnoyarsk, Russia

<sup>6</sup> Institute of Theoretical and Experimental Physics, Moscow

General Relativity rewritten in conformal units identifies conformal intervals with the real observational distances. This identification provides the basis for the explanation of all the epochs of the Universe evolution including Ia supernova luminosity long distance–redshift relation by the dominance of the Casimir vacuum energy of all physical fields. A set of arguments is discussed in favor of the fact that the SNe Ia data in the conformal units can be an evidence of the conformal twistor structure of the space-time as a nonlinear realization of the affine group, just like the nonlinear realization of chiral symmetry and phenomenological Lagrangian is an evidence of the quark structure of hadrons.

PACS: 95.30.Sf; 98.80.-k; 98.80.Es

### INTRODUCTION

The conformal symmetry as a basis for the construction of the General Relativity (GR) was independently introduced by Deser and Dirac [1,2]. In particular, Dirac formulated the conformal-invariant approach to the GR [2] as a new variational principle for the Hilbert action introducing a dilaton (scalar) field, in addition to the metric components  $g_{\mu\nu}$ .

The conformal treatment of gravity is supported by the Ogievetsky theorem [3] according to which the GR-diffeomorphism group  $\text{Diff } R^{(1,3)}$  can be obtained as the closure of two finite-dimensional groups: the 15-parameter conformal group and the 20-parameter affine group having the Poincaré group as a common subgroup. Further it was shown [4] that in the case of the dynamical affine symmetries the method of nonlinear realization of symmetry groups [5] leads to the Hilbert action of Einstein's gravitational theory expressed in terms

of the Fock simplex components [6]. The conformal invariance not only picks out the Einstein GR among several appropriated invariants of the nonlinearly realized affine symmetry, but it also establishes the conformal units which have been introduced before by several researchers including Dirac [2].

In the present paper, we discuss a set of observational data and facts that follow from the conformal units [1, 2] in comparison with the Einstein ones. In our conformal version of the GR (CGR), the conformal symmetry breaking happens due to the presence of the Casimir vacuum energy [7] in a finite volume of the Universe. In our approach, the Casimir vacuum energy substitutes the dark energy. It provides a good description of SNe Ia data [8] within the conformal cosmology [9, 10]. We found that the Universe horizon and the Planck least action postulate lead to the Planck scale hierarchy and the instance of the primordial particle creation from vacuum, with the Casimir vacuum energy being the source of the creation.

## 1. CONFORMAL GENERAL RELATIVITY

The Conformal General Relativity (CGR) is a nonlinear realization of the joint conformal and the affine  $A(4)$  symmetries in the factor space  $A(4)/L$  with the Lorentz subgroup  $L$  of the stable vacuum (here we use the concepts of the theory [5]). Recall that the affine group  $A(4)$  is the group of all linear transformations of the four-dimensional manifold  $x^\mu \rightarrow \tilde{x}^\mu = x^\mu + y^\mu + L_{[\mu\nu]}x^\nu + R_{\{\mu\nu\}}x^\nu$ , where  $y^\mu$  is a shift of coordinate and  $L_{[\mu\nu]}$  and  $R_{\{\mu\nu\}}$  are antisymmetric and symmetric matrices, respectively. A nonlinear realization of  $A(4)$  is based on finite transformations  $G = e^{iP \cdot x} e^{iR \cdot h}$  defined by means of the shift operator  $P$ , the proper affine transformation  $R$ , and the following Goldstone modes: four coordinates  $x_\mu$  and ten gravitational fields  $h$ . Further it was shown [4] that in the case of the dynamical affine symmetries the method of nonlinear realization of symmetry groups [5] leads to the Hilbert action of Einstein's gravitational theory expressed in terms of the Fock simplex components as  $W_E = -(M_{\text{Pl}}^2/16) \int d^4x \sqrt{-g} R^{(4)}$ , where  $R^{(4)}$  is the curvature. Taking into account the hidden conformal symmetry associated with a dilaton (scalar) field  $D$  and, consequently, transforming  $g_{\mu\nu}$  [2]

$$g_{\mu\nu} = \tilde{g}_{\mu\nu} e^{2D}, \quad (1)$$

we obtain the conformal-invariant action:

$$W_C \equiv -M_C^2 \frac{3}{8\pi} \int d^4x \left[ \frac{\sqrt{-\tilde{g}}}{6} R^{(4)}(\tilde{g}) e^{-2D} - e^{-D} \partial_\mu \left( \sqrt{-\tilde{g}} \tilde{g}^{\mu\nu} \partial_\nu e^{-D} \right) \right], \quad (2)$$

where  $M_C$  is a scale unit. It is defined in the Riemannian space-time where the conformal interval

$$ds_C^2 = \tilde{g}_{\mu\nu} dx^\mu dy^\nu = \tilde{\omega}_{(0)} \otimes \tilde{\omega}_{(0)} - \tilde{\omega}_{(b)} \otimes \tilde{\omega}_{(b)} \equiv \eta_{(\alpha)(\beta)} \tilde{\omega}_{(\alpha)} \otimes \tilde{\omega}_{(\beta)} \quad (3)$$

is identified with the measurable one, instead of the Einstein interval

$$ds_E^2 = g_{\mu\nu} dx^\mu dy^\nu. \quad (4)$$

If  $D = 0$ , one obtains  $W_C \equiv W_E$  and  $ds_C \equiv ds_E$ . Thus, the GR model based on the conformal and affine symmetry principles (described by the action (2)) differs from the original Einstein–Hilbert action  $W_E$  by the following elements and treatments. Namely:

1. Action (2) deals with the conformal geometrical interval (3)  $ds_C^2$  instead of the Einstein one (4)  $ds_E^2 = g_{\mu\nu} dx^\mu dx^\nu < ds_C^2$ .

2. The cosmological evolution in the CGR can be provided by the mean field dynamics of the dilaton zeroth mode instead of the homogeneous approximation [11] (see below).

3. The CGR contains the Newton coupling constant ( $G_N = M_C^{-2} e^{2D} = M_{\text{Pl}}^{-2}$ ) as the present-day value of the dilaton field  $D$ . We recall that the standard GR contains the effective Newtonian coupling constant as the absolute fundamental parameter of the equations of motion. In the CGR, the relation of the coupling constant to the Early Universe is clarified below.

In order to establish a relation between physical scales relevant for the Early Universe, we assume that there is a common source of the conformal symmetry breaking. We suppose that the Casimir vacuum energy of the Empty Universe could be naturally associated with this source (see below).

Hereafter, we use the *natural units*:

$$M_{\text{Pl}} \sqrt{3/(8\pi)} = c = \hbar = 1. \quad (5)$$

Taking into account Eqs. (3) and (4), the simplex components  $[\tilde{\omega}_{(0)}, \tilde{\omega}_{(b)}]$  can be written as

$$\tilde{\omega}_{(0)} = e^{-2D} N dx^0, \quad (6)$$

$$\tilde{\omega}_{(b)} = \mathbf{e}_{(b)i} dx^i + N_{(b)} dx^0, \quad (7)$$

where  $N_{(b)} = N^j \mathbf{e}_{j(b)}$  are the shift vector components, and  $N(x^0, x^j)$  is the lapse function. Here  $\tilde{\omega}_{(b)}$  are the linear forms defined via the triads  $\mathbf{e}_{(b)i}$  with a unit spatial metric determinant  $|\tilde{g}_{ij}^{(3)}| = 1$  known as the Lichnerowicz gauge [12]. This gauge fixes the scalar dilaton field  $D$  as the logarithm of the conformal factor:

$$D = -(1/6) \ln |g_{ij}^{(3)}|. \quad (8)$$

## 2. THE DILATON SCALAR FIELD

The group of invariance of the GR for the Dirac–ADM foliation is known as the kinematic subgroup of the general coordinate transformation [13]:

$$x^0 \rightarrow \tilde{x}^0 = \tilde{x}^0(x^0), \quad (9)$$

$$x^k \rightarrow \tilde{x}^k = \tilde{x}^k(x^0, x^1, x^2, x^3). \quad (10)$$

This group admits the decomposition of the dilaton into the sum of the zeroth and nonzerth harmonics:

$$D(x^0, x^1, x^2, x^3) = \langle D \rangle(x^0) + \overline{D}(x^0, x^1, x^2, x^3). \quad (11)$$

The introduction of the zeroth mode  $\langle D \rangle(x^0)$  is consistent with the Einstein cosmological principle of averaging of all scalar fields of the theory over a finite volume  $V_0 = \int_{V_0} d^3x$  [14] so that

$$\langle D \rangle(x^0) = V_0^{-1} \int_{V_0} d^3x D(x^0, x^1, x^2, x^3). \quad (12)$$

Note that the zeroth dilaton harmonics coincides by construction with the cosmological scale factor logarithm [11]

$$\langle D \rangle = -\ln a = \ln(1 + z). \quad (13)$$

Thus, in the finite volume  $V_0$  (taking into account Eqs. (11) and (13)), we have the following action:

$$W_C = \underbrace{W_{\text{Universe}}}_{=0 \text{ for } V_0=\infty} + W_{\text{graviton}} + W_{\text{potential}}, \quad (14)$$

$$W_{\text{Universe}}[\langle D \rangle, N_0] = -V_0 \int_{\tau_I}^{\tau_0} \underbrace{dx^0 N_0}_{=d\tau} \left[ \left( \frac{d\langle D \rangle}{N_0 dx^0} \right)^2 + \rho_{\text{Cas}}(\langle D \rangle) \right], \quad (15)$$

$$W_{\text{graviton}} = \int d^4x \frac{N}{6} \left[ v_{(a)(b)} v_{(a)(b)} - e^{-4D} R^{(3)}(\mathbf{e}) \right], \quad (16)$$

$$W_{\text{potential}} = \int d^4x N \left[ -v_D^2 - \underbrace{\frac{4}{3} e^{-7D/2} \Delta^{(3)} e^{-D/2}}_{\text{Newtonian potentials}} \right], \quad (17)$$

where

$$v_{\overline{D}} = \frac{1}{N} [(\partial_0 - N^l \partial_l) \overline{D} + \partial_l N^l / 3], \quad (18)$$

$$v_{(a)(b)} = \frac{1}{N} \left[ \omega_{(a)(b)}^R (\partial_0 - N^l \partial_l) + \partial_{(a)} N_{(b)}^\perp + \partial_{(b)} N_{(a)}^\perp \right] \quad (19)$$

are the velocities of the metric components and fields,  $\Delta = \partial_i [e_{(a)}^i e_{(a)}^j \partial_j]$  is the Beltrami–Laplace operator, and  $R^{(3)}(\mathbf{e})$  is the three-dimensional spatial curvature expressed in terms of the triads  $e_{(a)i}$ . Here we have introduced in action (15) the additional term  $\rho_{\text{Cas}}(\langle D \rangle)$ . The introduction of the finite volume  $V_0 = \int_{V_0} d^3 x <$

$\infty$  creates a dimensional parameter; therefore, it breaks the conformal symmetry. According to the general wisdom [5], this breaking leads to the appearance of a Goldstone mode [15,16]. It is just the zeroth harmonic  $\langle D \rangle$  that cannot be defined in the infinite volume. Note, however, that the Hamiltonian dynamics governed by the equations of motion must obey the conformal symmetry (see below). We will show that this source could be associated with the Casimir energy of the Universe giving a nonzero density contribution  $\rho_{\text{Cas}}(\langle D \rangle) \neq 0$ .

The choice of the zeroth dilaton mode  $\langle D \rangle$  as an evolution parameter has two consequences in the Hamiltonian approach. First, the zeroth dilaton mode canonical momentum density

$$P_{\langle D \rangle} = \frac{2}{V_0} \int_{V_0} d^3 x \sqrt{-g} g^{00} \frac{d}{dx^0} \langle D \rangle \equiv 2 \frac{d}{d\tau} \langle D \rangle = 2v_{\langle D \rangle} = \text{const} \neq 0 \quad (20)$$

can be treated as a generator of the Hamiltonian evolution in the field space of events [17,18]. We stress that the scale-invariance ( $D \rightarrow D + \Omega$ ) admits only a constant  $P_{\langle D \rangle}$ . In virtue of Eqs.(11) and (12), the Dirac Hamiltonian theory provides the orthogonality condition

$$\int_{V_0} d^3 x \overline{D}(x^0, x^1, x^2, x^3) \equiv 0. \quad (21)$$

This condition enables us to consider the zeroth and nonzerth components as independent ones.

The second consequence of the orthogonality condition (21) is that the nonzerth harmonics  $\overline{D}(x^0, x^1, x^2, x^3)$  do not depend on the evolution parameter. Therefore, one can consider these components as the gravitational Newton-type potentials due to the condition for the canonical momentum of the dilaton nonzerth modes

$$P_{\overline{D}}/2 = v_{\overline{D}} = [(\partial_0 - N^l \partial_l) \overline{D} + \partial_l N^l / 3] / N = 0. \quad (22)$$

This result fixes the longitudinal shift vector component.

As a result, we have

$$\int d^3x v_{\langle D \rangle} \cdot v_{\overline{D}} = 0 \quad (23)$$

that follows from Eqs. (11), (12), and (21). The orthogonality conditions (21) and (23) preserve the definite metrics in the Hilbert space of states [16].

### 3. CONFORMAL CASIMIR ENERGY AND UNIVERSE HORIZON

Let us consider the Early Universe. We assume that at the instance of creation the world was empty and finite in size. Therefore its energy can be associated only with the quantum Casimir energy of all physical fields in the given space. We will treat all those fields as massless since  $m(a) \xrightarrow{a \rightarrow 0} 0$  in the Early Universe epoch.

The Casimir energy of a massless field  $f$

$$H_{\text{Cas}}^{(f)} = \sum_{\mathbf{k}} \frac{\sqrt{\mathbf{k}^2}}{2} = \frac{\tilde{\gamma}^{(f)}}{d_{\text{Cas}}(a)} \quad (24)$$

depends on the geometry, size  $d_{\text{Cas}}$ , topology, boundary conditions, and spin (in particular, for a sphere of diameter  $d_{\text{Cas}}$ , the number of  $\tilde{\gamma} \sim 0.1-0.03$ ) [16]. For simplicity, we assume that the Universe has a spherical volume limited by the horizon.

It is natural to suggest that the energy of a massless field is proportional to the inverse visual size of the Universe  $d_{\text{Cas}}(a)$ . Assuming the same dependence for all fields, we define the total Casimir energy density of the Universe summing over all fields

$$\rho_{\text{t}} = \sum_f \frac{H_{\text{Cas}}^{(f)}}{V_0} = \frac{C_0}{d_{\text{Cas}}(a)}. \quad (25)$$

The key assumption of our model is that the Casimir dimension  $d_{\text{Cas}}(a)$  is equal to the Universe visual size (its horizon)

$$d_{\text{Cas}}(a) \equiv d_{\text{hor}}(a) = 2C_0^{-1/2} \int_{a_I \rightarrow 0}^a d\bar{a} d_{\text{Cas}}^{1/2}. \quad (26)$$

Equation (26) has the solution

$$d_{\text{Cas}}^{1/2}(a) = [C_0]^{-1/2} a \rightarrow d_{\text{Cas}}(a) = \frac{a^2}{C_0}. \quad (27)$$

Comparing Eq. (27) with the horizon

$$d_{\text{hor}}(a) = \frac{a^2}{H_0}, \quad (28)$$

one obtains

$$C_0 = H_0. \quad (29)$$

Thus, in our approach, parameter  $C_0$  is equal to the Hubble parameter  $H_0$ , which can be determined from the observations.

#### 4. HIERARCHY OF COSMOLOGICAL SCALES

Let us consider the Early Universe at the rigid state horizon (28). A hypothetical observer measures the conformal horizon  $d_{\text{hor}} = 2r_{\text{hor}}(z)$  as the distance that a photon covers within its light cone. The latter is determined by the zero interval equation  $d\eta^2 - dr^2 = 0$  during the photon lifetime in the homogeneous Universe, which is the subject of the condition  $\eta_{\text{hor}} = r_{\text{hor}}(z) = 1/[2H_0(1+z)^2]$ , in accordance with Eq. (27). This means that the four-dimensional space-time volume restricted by the horizon is equal to

$$V_{\text{hor}}^{(4)} = \frac{4\pi}{3} r_{\text{hor}}^3(z) \eta_{\text{hor}}(z) = \frac{4\pi}{3 \cdot 16H_0^4(1+z)^8}. \quad (30)$$

It is natural to assume that at the instance of the Universe origin the world was essentially quantum. Therefore, we claim that action (15) is the subject of the *Planck's least action postulate* so that

$$W_{\text{Universe}} = \rho_{\text{cr}} V_{\text{hor}}^{(4)}(a_{\text{Pl}}) = \frac{M_{\text{Pl}}^2}{H_0^2} \frac{(1+z_{\text{Pl}})^{-8}}{32} = 2\pi. \quad (31)$$

Using the present-day ( $\tau = \tau_0$ ) observational data for the Planck mass and the Hubble parameter at  $h \simeq 0.7$

$$\begin{aligned} M_C e^{\langle D \rangle(\tau_0)} &= M_{\text{Pl}} = 1.2211 \cdot 10^{19} \text{ GeV}, & \langle D \rangle(\tau_0) &= 0, \\ \frac{d}{d\tau} \langle D \rangle(\tau_0) &= H_0 = 2.1332 \cdot 10^{-42} \text{ GeV} \cdot h = 1.4332 \cdot 10^{-42} \text{ GeV}, \end{aligned} \quad (32)$$

we obtain from (31) the primordial redshift value

$$a_{\text{Pl}}^{-1} = (1+z_{\text{Pl}}) \approx \left[ \frac{M_{\text{Pl}}}{H_0} \right]^{1/4} \left[ \frac{4}{\pi} \right]^{1/8} / 2 \simeq 0.85 \cdot 10^{15}. \quad (33)$$

In other words, the Planck mass and the present-day Hubble parameter value (the main cosmological scales) are related to each other by the age of the Universe expressed in terms of the cosmological scale factor.

In field theories, characteristic scales associated with physical states are classified according to the Poincaré group representation [19]. In our approach, the Poincaré classification of energies arises from the decomposition of the mean particle energy  $\omega_\tau = a^2 \sqrt{\mathbf{k}^2 + a^2 M_0^2}$  conjugated to the dilaton time interval. We express this decomposition in the form

$$\langle \omega \rangle^{(n)}(a) = \left( \frac{a}{a_{\text{Pl}}} \right)^{(n)} H_0, \quad (34)$$

based on the primordial redshift value (33). This equation enables one to introduce the conformal weights  $n = 0, 2, 3, 4$  which correspond to: the dilaton velocity  $v_D = H_0$ , the massless energy  $a^2 \sqrt{\mathbf{k}^2}$ , the massive one  $M_0 a^3$ , and the Newtonian coupling constant  $M_{\text{Pl}} a^4$  (31), respectively. One can also include in these classifications the scale of the nonrelativistic particle  $H_0 = a_{\text{Pl}}^{-1} \cdot 10^{-13} \text{ cm}^{-1}$  with the unit conformal weight of its energy  $\omega_\tau^{\text{nonrel}} = a^1 \mathbf{k}^2 / M_0$ . As a result, the redshift leads to a hierarchy law of the present-day ( $a = 1$ ) cosmological scales

$$\omega_0^{(n)} \equiv \langle \omega \rangle^{(n)}(a) \Big|_{(a=1)} = \left( \frac{1}{a_{\text{Pl}}} \right)^{(n)} H_0, \quad (35)$$

shown in the Table.

**The hierarchy law of the cosmological scales in GeV** ( $M_{\text{Pl}}^* = \sqrt{3/(8\pi)} M_{\text{Pl}}$ )

$n = 0$	$n = 1$	$n = 2$	$n = 3$	$n = 4$
$H_0 \simeq 1.4 \cdot 10^{-42}$	$R_{\text{Cel.S}}^{-1} \simeq 1.2 \cdot 10^{-27}$	$k_{\text{0CMB}} \simeq 10^{-12}$	$\phi_0 \simeq 3 \cdot 10^2$	$M_{\text{Pl}}^* \simeq 4 \cdot 10^{18}$

The Table contains the scales corresponding to the Hubble parameter ( $n = 0$ ), the celestial system size ( $n = 1$ ), the cosmic microwave background mean wave-momentum ( $n = 2$ ), the electroweak scale of the SM ( $n = 3$ ), and the Planck mass ( $n = 4$ ). We conclude that the observational data testify that the cosmic evolution (34) of all these mean energies with conformal weights ( $n = 0, 1, 2, 3, 4$ ) has a common origin, which could be associated with the Casimir vacuum energy (see [22]).

## 5. SNE IA DATA AS THE EVIDENCES OF LONG CONFORMAL UNITS


A particular conformal cosmological model, based on the ideas discussed above, has been developed in [9, 10, 20, 21]. It was shown that the model leads to



a viable cosmology being in agreement with observations. For example, a good description of the modern supernovae type Ia (SNe Ia) data was constructed [9,10]. In the present paper, we show that the Casimir vacuum effect in a finite-size Universe could provide both the scale invariance breaking and the rigid state dominance, required in our model to describe the SNe Ia data.

Since the end of the last century, distant supernovae data is a widespread test for all the theoretical cosmological models in spite of the fact that the correctness of the hypothesis about SNe Ia as the perfect standard candles is still not proven [23]. Conformal cosmological models [24–26], where all observables are identified with the scale-invariant quantities of the GR introduced by Lichnerowicz [12], are also discussed among other possibilities [27].

Assuming that the supernovae type Ia are standard candles, one could use them to test cosmological theories. The Hubble Space Telescope team analyzed 186 SNe Ia [28] to test the Standard Cosmological model (SC) associated with expanded lengths in the Universe and evaluated its parameters (see the figure). We use the same sample to determine parameters of the Conformal Cosmological model (CC) with relative reference units of intervals, so that conformal quantities



$\mu(z)$ -dependence for cosmological models in SC and CC. The data points include 186 SNe Ia (the «gold» and «silver» samples) used by the cosmological supernova HST team. For a reference, we use the best fit for the flat standard cosmology model with  $\Omega_m = 0.27, \Omega_\Lambda = 0.73$  (the thick dashed line); the best fit for CC is shown with the thick solid line. For this CC model, we do not put any constraints on  $\Omega_m$

of the General Relativity are interpreted as observables. We concluded that the test is extremely useful and allows one to evaluate parameters of the model. From a formal statistical point of view, the best fit of the CC model is almost the same quality approximation as the best fit of the SC model with  $\Omega_\Lambda = 0.72$ ,  $\Omega_m = 0.28$ . As it was noted earlier, for CC models, a rigid matter component could substitute the  $\Lambda$ -term (or quintessence) existing in the SC model. We note that a free massless scalar field can generate such a rigid matter. We describe the results of our analysis for more recent «gold» data (for 192 SNe Ia).

## 6. SUMMARY

Any science is based on information. The units of information are a bit (1,0). The units of the quantum information are a 2-dimensional twistor  $(\cos \theta, \sin \theta e^{i\delta})$  as a fundamental representation of the conformal (C) group in the Penrose twistor program [29]. In accordance with this twistor program, the four-parametric matrix of the space-time coordinates  $\hat{x}_{A\dot{A}} = x_0 \hat{I}_{A\dot{A}} - x_j \sigma_{A\dot{A}}^j$  is constructed from the 2-dimensional twistor fundamental representations, like mesons (as a joint representation of  $SU(2)$ ) are constructed from the two-dimensional quark fundamental representation of  $SU(2)$ .

The next step in this analogy with the hadron physics (that is beyond the Penrose twistor program) is a nonlinear realization of the affine and conformal symmetries  $A(4) \otimes C$  in the factor-space  $K = A(4)/L$  in conformal units. This step is just like nonlinear realizations of the finite-parameter  $SU(2) \times SU(2)$  group over the vacuum stability subgroup  $SU(2)$ . Recall that this step leads to the effective chiral hadron Lagrangians constructed via the Maurer–Cartan linear forms without any reference to the underlying QCD theory. In this analogy, the Planck mass  $M_{Pl}$  plays the role of the weak decay constant parameter  $F_\pi \simeq 93$  MeV as a specific scale of hadron low-energy physics. One can see that the twistor dissociation (like QCD parton-type deep-inelastic scattering processes) can happen in the quantum theory of space-time when energy is greater than the Planck mass  $M_{Pl}$ . The question is what is the QCD analogy of such a twistor dissociation?

This hadron-like chain of the quantum theory of space-time proves that the supersymmetric unification can be based on the finite-parameter geometrization of all interactions via nonlinear realization of this super-affine group  $A[(2_b + 2_f) \times x(2_b + 2_f)] = A[8_b + 8_f]$ , where the role of twistors as the fundamental representation of the conformal group can be played by the supertwistors proposed in [30], together with the commutation-relation algebra of operators «superspace» conformal transformations associated with these supertwistors. This algebra forms a super-affine group and its nonlinear realization as the  $8_b + 8_f$  space-time. According to the hadron analogy, one can obtain the nonlinear realization, where

supercurvature  $R^{8_b+8_f}$  is a basis of the unified supersymmetric theory. These programs are supported by the last results in paper [31], where the gravi-electroweak and strong interactions were obtained by the unification of an 8-dimensional theory by compactification of four extra space dimensions in the theory with the curvature  $R^{(8)}$ .

**Acknowledgements.** The authors would like to thank M. Bordag, S. Deser, D. Ebert, A. Efremov, V. Gershun, Yu. Ignatev, E. Lukierski, and A. Zheltukhin for the useful discussions. VNP and AB were supported in part by the Bogoliubov–Infeld program. AFZ is grateful to the JINR Directorate for support.

## REFERENCES

1. *Deser S.* Scale Invariance and Gravitational Coupling // *Ann. Phys.* 1970. V. 59. P. 248.
2. *Dirac P. A. M.* Long Range Forces and Broken Symmetries // *Proc. Roy. Soc. London A.* 1973. V. 333. P. 403.
3. *Ogievetsky V. I.* Infinite-Dimensional Algebra of General Covariance Group as the Closure of Finite-Dimensional Algebras of Conformal and Linear Groups // *Lett. Nuovo Cim.* 1973. V. 8. P. 988.
4. *Borisov A. B., Ogievetsky V. I.* Theory of Dynamical Affine and Conformal Symmetries as Gravity Theory // *Theor. Math. Phys.* 1975. V. 21. P. 1179 (*Teor. Mat. Fiz.* 1974. V. 21. P. 329).
5. *Coleman S. R., Wess J., Zumino B.* Structure of Phenomenological Lagrangians. 1 // *Phys. Rev.* 1969. V. 177. P. 2239;  
*Volkov D. V.* Phenomenological Lagrangians // *Part. Nucl.* 1973. V. 4. P. 3; Preprint ITF-69-73. Kiev, 1969.
6. *Fock V.* Geometrization of Dirac's Theory of the Electron // *Z. Phys.* 1929. V. 57. P. 261 (in German).
7. *Actor A. A.* Scalar Quantum Fields Confined by Rectangular Boundaries // *Fortsch. Phys.* 1995. V. 43. P. 141;  
*Bordag M. et al.* *Advances in the Casimir Effect.* N. Y.: Oxford Univ. Press Inc., 2009.
8. *Riess A. G. et al. (Supernova Search Team Collab.).* Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant // *Astron. J.* 1998. V. 116. P. 1009;  
*Astier P. et al. (The SNLS Collab.).* The Supernova Legacy Survey: Measurement of  $\Omega(m)$ ,  $\Omega(\lambda)$  and  $W$  from the First Year Data Set // *Astron. Astrophys.* 2006. V. 447. P. 31.
9. *Behnke D. et al.* Description of Supernova Data in Conformal Cosmology without Cosmological Constant // *Phys. Lett. B.* 2002. V. 530. P. 20.
10. *Zakharov A. F., Pervushin V. N.* Conformal Cosmological Model Parameters with Distant SNe Ia Data: «Gold» and «Silver» // *Intern. J. Mod. Phys. D.* 2010. V. 19. P. 1875.

11. *Friedmann A.* Über die Krümmung des Raumes // *Z. Phys.* 1922. V. 10. P. 377 (Gen. Rel. Grav. 1999. V. 31. P. 1991) (in German);  
Über die Möglichkeit einer Welt mit konstanter negativer Krümmung des Raumes // *Z. Phys.* 1924. V. 21. P. 306 (in German).
12. *Lichnerowicz A.* L'intégration des équations de la gravitation relativiste et le problème des  $n$  corps // *J. Math. Pures and Appl. B.* 1944. V. 37. P. 23 (in French);  
*York J. W.* Gravitational Degrees of Freedom and the Initial-Value Problem // *Phys. Rev. Lett.* 1971. V. 26. P. 1656;  
*Kuchar K.* A Bubble-Time Canonical Formalism for Geometrodynamics // *J. Math. Phys.* 1972. V. 13. P. 768.
13. *Zelmanov A. L.* Chronometric Invariants and Accompanying Coordinates in General Relativity // *Dokl. AN USSR.* 1956. V. 107. P. 315 (in Russian);  
Kinematic Invariants and Their Relation to the Chronometric Invariants of Einstein's Theory of Gravity // *Dokl. AN USSR.* 1973. V. 209. P. 822 (in Russian).
14. *Einstein A.* Cosmological Considerations in the General Theory of Relativity // *Sitzungsber. Preuss. Akad. Wiss. Berlin*, 1917. P. 142.
15. *Goldstone J.* Field Theories with Superconductor Solutions // *Nuovo Cim.* 1961. V. 19. P. 154.
16. *Grib A. A., Mamaev S. G., Mostepanenko V. M.* Quantum Effects in Strong External Fields. St. Petersburg: Friedmann Lab. Publ., 1994.
17. *DeWitt B. S.* Quantum Theory of Gravity. 1: The Canonical Theory // *Phys. Rev.* 1967. V. 160. P. 1113.
18. *Wheeler J. A.* Lectures in Mathematics and Physics // *Batelle Rencontres: 1967 / Ed. by C. DeWitt and J. A. Wheeler.* N. Y., 1968.
19. *Wigner E. P.* On Unitary Representations of the Inhomogeneous Lorentz Group // *Ann. Math.* 1939. V. 40. P. 149.
20. *Barbashov B. M. et al.* Hamiltonian Cosmological Perturbation Theory // *Phys. Lett. B.* 2006. V. 633. P. 458.
21. *Arbuzov A. B. et al.* Conformal Hamiltonian Dynamics of General Relativity // *Phys. Lett. B.* 2010. V. 691. P. 230.
22. *Pervushin V. N. et al.* Conformal and Affine Hamiltonian Dynamics of General Relativity. arXiv:1109.2789 [gr-qc].
23. *Panagia N.* High Redshift Supernovae: Cosmological Implications // *Nuovo Cim. B.* 2005. V. 120. P. 667.
24. *Behnke D. et al.* Description of Supernova Data in Conformal Cosmology without Cosmological Constant // *Phys. Lett. B.* 2002. V. 530. P. 20.
25. *Behnke D.* Conformal Cosmology Approach to the Problem of Dark Matter. PhD Thesis. Rostock Report MPG-VT-UR 248/04. 2004.
26. *Blaschke D. B. et al.* Cosmological Production of Vector Bosons and Cosmic Microwave Background Radiation // *Phys. At. Nucl.* 2004. V. 67. P. 1050;

- 
- Barbashov B. M. et al.* Hamiltonian General Relativity in Finite Space and Cosmological Potential Perturbations // Intern. J. Mod. Phys. A. 2006. V. 12. P. 5957;  
*Barbashov B. M. et al.* The Hamiltonian Approach to General Relativity and CMB Primordial Spectrum // Intern. J. Geom. Meth. Mod. Phys. 2007. V. 4. P. 171.
27. *Riess A. G. et al. (Supernova Search Team Collab.)*. The Farthest Known Supernova: Support for an Accelerating Universe and a Glimpse of the Epoch of Deceleration // Astrophys. J. 2001. V. 560. P. 49;  
*Tegmark M.* Measuring the Metric: A Parametrized Post-Friedmanian Approach to the Cosmic Dark Energy Problem // Phys. Rev. D. 2002. V. 66. P. 103507.
28. *Riess A. G. et al.* Type Ia Supernova Discoveries at  $z > 1$  from the Hubble Space Telescope: Evidence for Past Deceleration and Constraints on Dark Energy Evolution // Astrophys. J. 2004. V. 607. P. 665.
29. *Penrose R.* Relativity, Groups and Topology. London: Gordon and Breach, 1964;  
*Chernikov N., Tagirov T.* Quantum Theory of Scalar Fields in de Sitter Space-Time // Ann. Inst. Henri Poincarè. 1968. V. 9. P. 109.
30. *Litov L. B., Pervushin V. N.* Quantum Supertwistors and Fundamental Superspaces // Phys. Lett. B. 1984. V. 147. P. 76.
31. *Vladimirov Yu. S., Gubanov A. N.* Unification of Gravi-Electroweak and Strong Interactions in the 8-Dimensional Theory // Grav. Cosm. 1999. V. 5. P. 277.