

The State Research Center of Russian Federation
BUDKER INSTITUTE OF NUCLEAR PHYSICS

A.M. Kudryavtsev, K.V. Lotov, and A.N. Skrinsky

ON ENERGY LOSS OF NARROW AND DENSE
ULTRARELATIVISTIC BUNCH IN PLASMA

Budker INP 96-64

NOVOSIBIRSK

1996

On energy loss of narrow and dense ultrarelativistic bunch in plasma

A.M. Kudryavtsev, K.V. Lotov, and A.N. Skrinsky

Budker Institute of Nuclear Physics SB RAS
630090 Novosibirsk, Russia

Abstract

Energy balance of a particle bunch propagating through the plasma is considered for the bunches of a high density (compared to plasma) and a small radius (compared to anomalous skin depth). Energy loss to scattering of near-axis plasma electrons is estimated. This component of energy balance is shown to be important but not excessively high for the parameters interesting for high energy physics. In particular, the energy loss to near-axis electrons is proved to remain finite as the bunch radius tends to zero.

Energy balance of a particle bunch propagating through the plasma is one of the key questions for plasma-based advanced accelerator techniques [1, 2]. Earlier [3], this problem was studied for the bunches of a low density (as compared to that of the plasma) with the most attention paid to beam loading issues. In this paper we consider the energy balance of a dense (compared to plasma) and narrow (compared to the anomalous skin depth c/ω_p) axisymmetric bunch. In particular, we show that the energy transferred from the bunch to near-axis plasma electrons remains finite as the bunch radius tends to zero (while the bunch length remains finite).

The latter issue is especially important for potential applications of plasma to linear collider for TeVs energy range. In this case the accelerated beams have to be of extremely small emittances and radii to provide acceptable luminosity. The bunch population remains much smaller than the number of plasma electrons per $(c/\omega_p)^3$, but the bunch density is much higher than that of the plasma (n_0).

The total energy balance of such a bunch is composed of two parts: the energy loss due to acceleration of near-axis plasma electrons, and the energy extracted from (or transferred to) remote plasma layers (at radii $\sim c/\omega_p$). The latter does not contain any peculiar to narrow bunch features. It can be easily found in particle-in-cell [3] or finite-difference [4] simulations via calculation of the longitudinal electric field. In the nonlinear plasma wave (with the electric field amplitude of the order of $\sqrt{4\pi n_0 mc^2}$), this long-range interaction can provide the energy gain up to $m c \omega_p$ per particle per unit length, where m is the electron mass.

The energy lost to near-axis electrons can not be calculated so straightforwardly. Fluid codes fail to solve this problem because of essentially non-hydrodynamical motion of plasma electrons near the dense bunch. The correct value of energy losses can, in principle, be obtained with existing particle-in-cell codes, but introduction of the additional length-scale (associated with the bunch radius) greatly increases the number of particles required and, thus, prohibits the study of extremely narrow bunches. The aim of this paper is to give correct estimate of this non-hydrodynamic energy loss for very narrow bunches of a very high density. Note that for this estimate it is not necessary to specify the profile of the plasma wave accelerating the bunch. Since the energy acquired by near-axis plasma

electrons is much greater than their initial energy (in the plasma wave), the wave profile affects the energy loss only through the change of plasma electron density.




Figure 1: The energy loss for electron (e) and positron (p) bunches.

To establish the upper limit of energy losses, we consider an infinitely narrow bunch (composed of N_b particles) propagating through the unperturbed plasma of density n_0 . The particles are assumed to be uniformly distributed over the bunch length l . Even for zero radius bunch, the energy loss per unit length (W) turns out to be finite:

$$W = N_b \frac{mc^2}{l} F(L). \quad (1)$$

Here

$$L = \frac{2l\omega_p}{c} \sqrt{\frac{N_b e^2}{mc^2 l}}, \quad (2)$$

$\omega_p = \sqrt{4\pi n_0 e^2 / m}$ is the plasma electron frequency, other notation is conventional. The dimensionless functions $F(L)$ for negatively and positively charged bunches are shown in Fig. 1 by solid lines. Similar functions for the bunches of a finite width (of the radius c/ω_p , with the particles being uniformly distributed over the bunch cross-section) are shown by dots. The difference between these graphs represents the energy loss to the near-axis plasma layer. It is seen that the energy loss of the narrow bunch is several times higher than that of a “wide” bunch.

Expression (1) and the functions $F(L)$ are obtained as follows. The radial electric field E_r and the azimuthal magnetic field H_φ of the infinitely narrow ultrarelativistic bunch may be approximated by the formula

$$E_r(r) = H_\varphi(r) = \frac{2eN_b}{rl} \exp\left(-\frac{r\omega_p}{c}\right) \quad (3)$$

(we use cylindrical coordinates with the z -axis being the direction of bunch propagation). This approximation is good for $r \ll c/\omega_p$, while for $r \gtrsim c/\omega_p$ the screening of the bunch field by the plasma (see, e.g., [5]) is taken into account only qualitatively through the exponential factor (fully self-consistent account of this region unnecessarily complicates the calculation of energy loss to near-axis layer). In particular, we neglect temporal

changes of the fields there caused by the bunch-driven displacement of plasma electrons. It is self-evident that the bunch can be treated as “rigid” and unaffected by the fields appearing in plasma.

Within the considered model, the energy of plasma electrons and the z -component of their generalized momentum are conserved:

$$\gamma mc^2 - e\Phi = \text{const}, \quad \gamma mv_z - \frac{e}{c}\Phi = \text{const}, \quad (4)$$

where $\Phi(r)$ is the electrostatic potential and the z -component of vector potential at the same time:

$$E_r(r) = H_\varphi(r) = -\frac{\partial\Phi}{\partial r}, \quad (5)$$

γ is the relativistic factor of plasma electrons, and e is the elementary charge ($e > 0$). The total energy loss is

$$W = n_0 \int_0^\infty e(\Phi(r_f) - \Phi(r_0)) 2\pi r_0 dr_0, \quad (6)$$

where r_0 corresponds to the initial position of a plasma electron (before the bunch disturbs the plasma), and r_f is its final radius (after the bunch passage).

For the electron bunch, we can find r_f for any specified r_0 by mere equating the electron travel time and the duration of field action:

$$\int_{r_0}^{r_f} \frac{dr}{v_r} = \frac{l}{c} + \frac{1}{c} \int_{r_0}^{r_f} \frac{v_z}{v_r} dr \quad (7)$$

(the integral in the right-hand side of (7) represents the distance travelled by the electron in z -direction). Since, from (4),

$$v_r = c \frac{\sqrt{2\Delta\Phi}}{1 + \Delta\Phi}, \quad v_z = c \frac{\Delta\Phi}{1 + \Delta\Phi}, \quad \text{where } \Delta\Phi = \frac{e(\Phi(r) - \Phi(r_0))}{mc^2}, \quad (8)$$

we can rewrite (7) as

$$l = \int_{r_0}^{r_f} \frac{dr}{\sqrt{2\Delta\Phi}},$$

whence the dependence $r_f(r_0)$ can be easily found numerically. Then numerical integration of (6) yields the formula (1) and the function $F(L)$.

For the positively charged bunch, the energy loss can be found in a similar way with the little complication arising from the oscillatory motion of near-axis plasma electrons (Fig. 2). Energy loss for wide bunches is calculated analogously.

The finiteness of the energy loss can be proved analytically as well. For the electron bunch this is evident, since the energy acquired by a plasma electron may be majorized by the function $e(\Phi(\infty) - \Phi(r_0))$, which has the integrable logarithmic singularity at $r_0 = 0$.




Figure 2: The final radius of plasma electrons as the function of their initial radius for electron (e) and positron (p) bunches with $L = 0.1$.

For the positively charged bunch the situation is more complicated, since the infinite energy is acquired by electrons for which

$$\int_0^{r_0} \frac{dr}{\sqrt{2\Delta\Phi}} = \frac{l}{2k+1} \quad (k = 0, 1, 2, \dots),$$

or

$$r_0 \approx \frac{c}{\omega_p} \cdot \frac{L}{(2k+1)G}, \quad G = \int_0^1 \frac{dx}{\sqrt{-\ln x}} = \sqrt{\pi} \quad (9)$$

(here we take into account the logarithmic behavior of Φ near the axis). The vicinity of each logarithmical singularity (9) makes a finite contribution to W , but there is an infinity of them. To majorize the total energy loss, we consider the contribution ΔW_k of the ring

$$r_k^0 = \frac{Lc}{2\sqrt{\pi}(k+1)\omega_p} < r_0 < r_k = \frac{Lc}{\sqrt{\pi}(2k+1)\omega_p}. \quad (10)$$

The final radius r_f is determined from the requirement

$$L = \frac{\sqrt{\pi}(2k+1)\omega_p}{c} r_0 + \int_0^{r_f} \frac{dr \cdot \omega_p/c}{\sqrt{\ln(r_0/r)}}, \quad (11)$$

whence

$$\frac{dr_f}{dr_0} = -\sqrt{\pi}(2k+1)\sqrt{\ln(r_0/r_f)}.$$

Then

$$\begin{aligned} \Delta W_k &\approx \frac{4\pi n_0 e^2 N_b}{l} \int_{r_k^0}^{r_k} r_0 \ln(r_0/r_f) dr_0 = -\frac{m\omega_p^2 N_b}{\sqrt{\pi} l (2k+1)} \int_{r_0=r_k^0}^{r_0=r_k} r_0 \sqrt{\ln(r_0/r_f)} dr_f, \\ \Delta W_k &< \frac{m\omega_p^2 N_b r_k^2}{\sqrt{\pi} l (2k+1)} \int_0^1 \sqrt{\ln(r_k/r_f)} \frac{dr_f}{r_k} = N_b m c^2 \frac{L^2}{2\pi l (2k+1)^3}. \end{aligned} \quad (12)$$

It is easy to show that the contribution of the ring $r_{k+1} < r_0 < r_k^0$ is less than ΔW_k . Since the series

$$\sum_{k=0}^{\infty} \frac{1}{(2k+1)^3}$$

converges, we conclude that the energy loss of the positively charged bunch is finite.

It is interesting to note why the calculation of energy loss in terms of “instant push” gives the erroneous result:

$$p \sim \frac{e|E_r|l}{c} \propto \frac{1}{r_0}, \quad W \propto \int 2\pi r_0 \cdot \frac{p^2}{2m} dr_0 \propto \int \frac{dr_0}{r_0} \quad (\text{diverges}). \quad (13)$$

This is because near-axis plasma electrons rapidly pass to the region of either smaller field (electron bunch) or opposite-sign field (positron bunch). As a result, the estimate (13) turns out to be unacceptably overstated.

The general conclusion of the paper is that the non-hydrodynamic energy losses of a dense bunch travelling in plasma are important but not excessively high for the parameters interesting for high energy physics.

References

- [1] **J. S. Wurtele** *The role of plasma in advanced accelerators.*— Phys. Fluids B, 1993, v. 5, №7, p. 2363–2370.
- [2] **E. Esarey, P. Sprangle, J. Krall, and A. Ting** *Overview of plasma-based accelerator concepts.* — IEEE Trans. Plasma Sci., 1996, v. 24, №2, p. 252–288.
- [3] **S. Wilks, T. Katsouleas, J. M. Dawson, P. Chen, and J. J. Su** *Beam loading in plasma waves.* — IEEE Trans. Plasma Sci., 1987, v. PS-15, №2, p. 210–217.
- [4] **B. N. Breizman, T. Tajima, D. L. Fisher, and P. Z. Chebotaev** *Excitation of Nonlinear Wake Field in a Plasma for Particle Acceleration.* — In: Research Trends in Physics: Coherent Radiation and Particle Acceleration, edited by A. Prokhorov, 1992, p. 263–287 (AIP Press, New York, 1992).
- [5] **R. Keinigs and M. E. Jones** *Two-dimensional dynamics of the plasma wakefield accelerator.* — Phys. Fluids, 1987, v. 30, №1, p. 252–263.

A.M. Kudryavtsev, K.V. Lotov, and A.N. Skrinsky

**On energy loss of narrow and dense
ultrarelativistic bunch in plasma**

A.M. Кудрявцев, К.В. Лотов, А.Н. Скрипский

**О потерях энергии узкого и плотного
ультрарелятивистского сгустка в плазме**

Budker INP 96-64

Ответственный за выпуск А.М. Кудрявцев

Работа поступила 23.09. 1996 г.

Сдано в набор 25.09.1996 г.

Подписано в печать 25.09.1996 г.

Формат бумаги 60×90 1/16 Объем 0.7 печ.л., 0.6 уч.-изд.л.

Тираж 200 экз. Бесплатно. Заказ № 64

Обработано на IBM PC и отпечатано на
ротапинтере ГНЦ РФ "ИЯФ им. Г.И. Будкера СО РАН",
Новосибирск, 630090, пр. академика Лаврентьева, 11.